

Original Research Article

Green and Efficient Synthesis of Baclofen

Pearl Rhea Fernandes¹, Ashwini Madnala², Nisha Donde³, Amit Pawar³, Maryappa sonawale*⁴

¹Department of Food Nutrition and Dietetics, Milagres College, Hampankatta Mangalore, India

²SIES College of Arts, Science and Commerce Sion, Mumbai, India

³Karmaveer Bhaurao Patil College, Vashi, Navi Mumbai, 400703, Maharashtra, India

⁴Veer wajekar ASC College Phunde, Mahalan Vibhag, Phunde, Uran Navi Mumbai Maharashtra India

ARTICLE INFO

Article history

Submitted: 2022-05-14

Revised: 2022-05-27

Accepted: 2022-06-10

Available online: 2022-06-22

Manuscript ID: AJCB-2205-1120

DOI: 10.22034/ajcb.2022.342298.1120

KEYWORDS

GABA agonist

4-Chlorobenzaldehyde

Cyanohydrin

Oxidizing agent

Pyridinium

Wittig type reaction.


Hydrolysis

Reduction

ABSTRACT

Baclofen is a Gamma Amino Butyric acid (GABA) agonist to relax skeletal muscles. It is especially effective in treating muscular spasticity. In this research, we now describe the synthesis of Baclofen using a new approach which begins with 4-chlorobenzaldehyde. We have carried all reactions by the green method. It gives the better yield. The identity of this product was confirmed by spectral analysis and compared with the standard Baclofen. New and eco-friendly approach was developed for the synthesis

GRAPHICAL ABSTRACT



* Corresponding author: Maryappa sonawale

✉ E-mail: maryappaconawale11@gmail.com

© 2022 by SPC (Sami Publishing Company)



1. Introduction

Baclofen is a gamma amino butyric acid (GABA) agonist for skeletal muscle relaxation. It is especially effective in treating spasticity. It interacts with both GABAA and GABAB receptors [1-3]. Mostly the GABA-A receptors are primarily found in the frontal lobe, GABA B receptors are more commonly found in the thalamus and the dorsal horn of the body's spinal cord [3-5]. Very fast synaptic suspension is mediated by GABAA receptors linked to chloride ion channels in the cells. GABA-B receptors are insensitive to bicuculline and linked to neuronal potassium and different calcium channels through G-proteins, allowing them to mediate delayed synaptic inhibition by raising potassium and lowering calcium conductance [6-7]. The antispasmodic and muscle relaxant baclofen, which is presently the sole selective and therapeutically relevant GABAB agonist, activates the GABAB receptors [8]. There are various accounts in the literature concerning the synthesis of Baclofen because of its biological and pharmacological value [9]. Although these publications employ diverse methodologies, the reagents used are costly [10-11], or the yield is poor. There are different methods used in the synthesis [12-18]. Various methods are used during the synthesis, including the preparation of various nanoparticles from plant materials [18-24]. In most cases, plant-derived nanoparticles are synthesized using phytochemical analysis [25-32]. A variety of heterocyclic compounds are synthesized using the Green Approach [33-35]. An alternate strategy to the synthesis of Baclofen is discussed in this work, which produces a high-yielding product and green pathway used for the synthesis. We have carried out the experiments by using green protocol, various new catalyst is used for reaction. We have reduced the cost of the production. All reaction gives above 80 % yield.

2. Experimental

All chemical components were reagent grade and analyzed using TLC on silica gel plates and U.V. light. Shimadzu IR 120 without KBR pellet was used to record I.R. spectra. Varian-400 spectrometer was used to record ¹H NMR spectra.

2.1. Procedure for the synthesis of pchlorobenzylidene-bis-acetoacetic ester (Step I)

A mixture of 2.8 gm of the para -chlorobenzaldehyde and 5.2 g of ethyl acetoacetate was treated at 0 °C with 4 mL of Morpholine as a base, kept it at 0-5 °C for 60-70 min. Then, it was sonicated at 20-25 °C for 12 hours. After that, 20.0 ml of ethanol was added slowly. The reaction mixture was completely cooled and filtered. The residue was washed with 70% ethanol until a solid white product was obtained.

2.2. Synthesis of β(chlorophenyl) glutaric acid derivative (Step II)

The solution was taken to have a temperature of 50-60 °C of 20.0 g of Lithium hydroxide in 15.0 mL water added to para -chlorobenzylidene-bisacetoacetic ester, which was prepared in step I. Then, this reaction mixture was subjected to sonication during this maintenance at 90-95 °C for almost 1 hour. After that, check the reaction progress using thin layer chromatography. After completing the reaction, the reaction mixture was diluted with two volumes of water. Wash the aqueous layer washed with ether. Then acidify the aqueous layer with 5.5 mL of concentrated hydrochloric acid and chilled thoroughly and filtered. The filter cake was washed several times with ice water and dried in the vacuum on a rot evaporator at 60 °C, then dry it properly.

2.3. Synthesis of β (p-chlorophenyl) glutarimid (Step III)

1.5 g. of β (p-chlorophenyl) glutaric acid was taken and synthesized in step II. Then, it was dissolved in 10.0 mL of distilled water. Next, 3.5 mL of concentrated ammonium hydroxide was added to this reaction mixture. After that, the color of the solution was removed. The solution was treated with charcoal and filtered over the

hyflo. Then, the mixture was heated in a proper open flask until the mixture microwave oven reached product formation and checked using the thin layer chromatography. Then after, 5 ml of the alcohol was added. The resulting reaction mixture was heated in the microwave until boiling. The completion was then diluted with 100 ml of hot water, stirred, completely cooled, and filtered. Finally, the filter cake was rinsed with ice water and dried in vacuum at 60 °C.




Fig. 1 Process for synthesis for balcofen

2.4. Procedure for the synthesis of 4-Amino-3-(4-chlorophenyl) butanoic acid (Baclofen) (Step IV)

0.5 g of the above product β (p-chlorophenyl) glutarimide was cooled to 10 °C to 15 °C in an ice bath. Next, to this reaction mixture, a solution of 0.5 g of sodium hydroxide in 2.0 mL of water was prepared, and then 0.4 g of bromine was added to the above reaction mixture over 20 minutes. The reaction mixture was then sonicated at 20-25 °C. for 4 hours. The reaction solution was then carefully adjusted to pH=7 with dilute hydrochloric acid (this is an important step). The microcrystalline 4-amino 3 (4-chlorophenyl) fatty acid (γ -amino β (p-

chlorophenyl) fatty acid) is then separated as a product. The product was washed with a suitable solvent.

3.Result and Observations

3.1.P-chlorobenzylidene-bis-acetoacetic ester

Yield:91%;

MP: 153-156 °C;

IR (Without KBr); ν 654, 828, 1470, 1580, 1720, 1740, 2960,3180, 3560, cm^{-1} ;

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 1.2 (2t, 6H), 1.5 (s, 3H), 2.9 (m,3H), 3.8 (m, 4H), 7.28 (q, 4H).

3.2. Synthesis of β (p-chlorophenyl) glutaric acid

Yield: 90%;

MP: 165-169°C;

IR (without KBr); ν 648, 828, 968, 1498, 1590, 1715, 2480-3420, cm^{-1} ;

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 2.7 (m, 4H), 3.4 (m, 1H), 7.5 (m, 4H), 12 (s, 2H).

3.3. β (p-chlorophenyl) glutarimide

Yield: 88%;

MP: 128-129°C;

IR (without KBr); ν 640, 820, 960, 1490, 1600, 1710, 2400-3400, cm^{-1} ;

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ 2.7 (m, 4H), 3.4 (m, 1H), 7.5 (m, 4H), 12 (s, 2H).

3.4. 4-Amino-3-(4-chlorophenyl) butanoic acid (baclofen)

Yield: 88%;

MP: 208-209°C;

IR (without KBr); 1098, 1498, 968, 1538, 1610, 1585, cm^{-1} .

4. Results and Discussion

We performed all the reactions on the Greenway. Various condensation reactions were used such as B. Claisen condensation. It is condensed to form a cyclic imide, and finally forms a Hofmann rearrangement. All three reactions can follow the general procedure used for the Green Protocol. In the reaction of ethyl acetoacetate with chlorobenzaldehyde, it is used in the alkoxide sodium / reaction to make ethyl acetoacetate an active nucleophile or to form a nucleophile that excels in the reaction required for Claisen condensation. Strong bases such as other bases are required. The primary medium can be further used for hydrolyzing ester molecules very easily to form acid molecules. Finally, the reaction of (Pchlorophenyl) glutaric acid with ammonium hydroxide or other bases can be used to produce (Pchlorophenyl) glutarimide. Next, in the Hofmann rearrangement, the reaction between bromine

and sodium hydroxide is converted into an imide, carbon dioxide is removed after hydrolysis, and baclofen is used as the final API.

5. Conclusion

To sum up, the synthesis of the different baclofen derivatives is in the final step. This study proposes a 4-step green, the cost-effective, straightforward, and synthetic pathway that has the potential for large-scale drug creation in the industrial synthesis of the product. This green strategy is used for Baclofen synthesis. During this, the synthetic approach uses inexpensive chemicals and leads to an overall yield of around 80% along with different green protocols.

References

- [1] Ogunkemi Risikat Agbeke Iyun, Ahmed Jibrin Uttu, Muhammad Sani Sallau, Hamisu Ibrahim. GC-MS Analysis of Methanol Extract of *Strychnos Innocua* (Delile) Root Bark. *Adv. J. Chem. A*, 2022, 5(2), 104-117
- [2] Adebayo Tajudeen Bale, Tolulope Mojisola Fasina, Rafiu Olarewaju Shaibu. Synthesis and Biological Study of Substituted 2''-Hydoxy, 2'',4''-dichloro Chalcones and Their Co(II), Cu(II) and Ni(II) Complexes for Their Antioxidant and Antimicrobial Potentials. *Adv. J. Chem. A*, 2022, 5(2), 94-103
- [3] Samuel Ndaghiya Adawara, Gideon Adamu Shallangwa, Paul Andrew Mamza, Ibrahim Abdulkadir. In-silico Design of Oxadiazole Hybrids as Potential Inhibitors of Dengue Virus NS2B-NS3 Protease. *Adv. J. Chem. A*, 2022, 5(2), 118-137.
- [4] D.A. Williams, T.L. Lemke, *Foye's Principles of Medicinal Chemistry, 5th Ed.* Lippincott Williams & Wilings, Philadelphia, (2002) 492-503.
- [5] Ifeoluwa Israel Alao, Ifeoluwa Peter Oyekunle, Kingsley O. Iwuozor, Ebuka Chizitere Emenike, Green synthesis of Copper Nanoparticles and

- Investigation of its Antimicrobial Properties, Ad. J. Chem. B, 4 (2022) 39-52.
- [6] Olaniyi O. Bashir, Oluwaniyi O. Omolara, Oloruntele O. Ibrahim, Sekoni A. Hauwa, Investigation of physicochemical and fatty acid composition of oils from ripe and unripe Blighia sapida fruit, Ad. J. Chem. B, 4 (2022) 53-61
- [7] Ifeoluwa Israel Alao, Ifeoluwa Peter Oyekunle, Kingsley O. Iwuozor, Ebuka Chizitere Emenike, Green synthesis of Copper Nanoparticles and Investigation of its Antimicrobial Properties, Ad. J. Chem. B, 4 (2022) 39-52.
- [8] Masood Bagheri Sadr, Kim Lo Han. Investigation of standard natural gas synthesis technologies, Ad. J. Chem. B, 4 (2022) 62-74.
- [9] N.A. Mirgane, V.S. Shivankar, S.B. Kotwal, G.C. Wadhawa, M.C. Sonawale, Degradation of dyes using biologically synthesized zinc oxide nanoparticles, *Materials Today: Proceedings*, 37(2021) 849-853. <https://doi.org/10.1016/j.matpr.2020.06.037>.
- [10] N.A. Mirgane, V.S. Shivankar, S.B. Kotwal, G.C. Wadhawa, M.C. Sonawale, the Waste pericarp of ananas comosus in green synthesis zinc oxide nanoparticles and their application in wastewater treatment, *Materials Today: Proceedings*, 37 (2021) 886-889. <https://doi.org/10.1016/j.matpr.2020.06.045>.
- [11] S.S. Nayak, N.A. Mirgane, V.S. Shivankar, K.B. Pathade, G.C. Wadhawa, Adsorption of methylene blue dye over activated charcoal from the fruit peel of plant hydnocarpus pentandra, *Materials Today: Proceedings*, 37 (2021) 2302-2305. <https://doi.org/10.1016/j.matpr.2020.07.728>.
- [12] D.D. Patil, K.D. Mhaske, C.G. Wadhawa, Antibacterial and Antioxidant study of Ocimum basilicum Labiatae (sweet basil), *Journal of Advanced Pharmacy Education & Research*, 2 (2011) 104-112.
- [13] P.D. Dinanath, W.C. Gurumeet, Antibacterial, antioxidant and anti-inflammatory studies of leaves and roots of Solanum xanthocarpum. *Unique J Ayurvedic Herb Med.*, 3(2013) 59-63.
- [14] D.K. Mhaske, D.D. Patil, G.C. Wadhawa. Antibacterial activity of methanolic extract from rhizome and roots of Valerianawallichii. *International Journal on Pharmaceutical and Biomedical Research*, 2(2011)107-111.
- [15] D.D. Patil, D.K. Mhaske, M.P. Gurumeet, C. Wadhawa, Antibacterial and antioxidant, anti-inflammatory study of leaves and bark of Cassia fistula. *Int J Pharm.*, 2(2012)401-405.
- [16] G.C. Wadhawa, M.A. Patare, D.D. Patil, D.K. Mhaske, Antibacterial, antioxidant and anti-inflammatory studies of leaves and roots of Anthocephalus kadamba. *Universal Journal of Pharmacy*, 2013.
- [17] S.S. Nayak, N.A. Mirgane, V.S. Shivankar, K.B. Pathade, G.C. Wadhawa, Degradation of the industrial dye using the nanoparticles synthesized from flowers of plant Ceropogia attenuata, *Materials Today: Proceedings*, 37(2021) 2427-2431. <https://doi.org/10.1016/j.matpr.2020.08.274>.
- [18] AbdulRazaq Tukur, Nasiru Malam Musa, Jimoh Tajuddeen Abdullahi, James D Habila and Martha Omotola Bamidele, Phytochemical Screening and the Effect of Methanolic Leaves Extract of Senna mimosoides on Inflammatory Stimulus Induced Leukocyte Mobilization (In-vivo), Ad. J. Chem. B, 4 (2022) 29-38
- [19] D.D. Patil, D.K. Mhaske, M.A. Patare, G.C. Wadhawa, Antibacterial and antioxidant, anti-inflammatory study of leaves and root Delonix regia. *International Journal of chemical and pharmaceutical sciences*, 3(2014)131-5
- [20] D. D. Patil, D. K. Mhaske, M.A. Patare, G.C. Wadhawa, Antibacterial and antioxidant, anti-inflammatory study of leaves and root Delonix regia. *International Journal of chemical and pharmaceutical sciences*, 3(2014) 131-5
- [21] A.K. Valvi, S.S. Nayak, V.S. Shivankar, G.C. Wadhawa, Nano catalyst derived from plant

- Polyzygus Tuberosus: A suitable for efficient and green synthesis of benzimidazole derivatives, *Materials Today: Proceedings*, 50(2022) 1522-1526. <https://doi.org/10.1016/j.matpr.2021.09.102>.
- [22] D.D.Patil, A.K.Deshmukh, G.C.Wadhawa, Diuretic activity of root and bark of erythrina indica lam. *International Journal of Pharmaceutical Sciences and Research.*, 2 (2011) 1811-1813
- [23] S.S. Nayak, N.A. Mirgane, K.B. Pathade, V.S. Shivankar, G.C. Wadhawa. Phytochemical analysis, antioxidant and anti-inflammatory activity of leaves and bark of Ceropogon rollae Hemadri, *Plant Sci. Today*, 8 (2021) 425-428. <https://doi.org/10.14719/pst.2021.8.3.906>
- [24] P.S. Gaikar, V.S. Shivankar, P.A.Patil, A.U.Chavan, G.C.Wadhawa, Preliminary Phytochemical Analysis and Antioxidant, Anti-Inflammatory Activity Of Dicliptera Ghatica Santapau. *Int. J. of Aquatic Science*, 12(2021) 4973-4980.
- [25] N.A. Mirgane, A. Chandore, V. Shivankar, Y. Gaikwad, G.C. Wadhawa, Phytochemical Study and Screening of Antioxidant, Anti-inflammatory Typhonium Flagelliforme, *Res. J. Pharm. Technol.*, 14(2021) 2686-2690.
- [26] D.D. Patil, G.C. Wadhawa, A.K. Deshmukh. One Pot Synthesis of Nitriles from Aldehydes and Hydroxylamine Hydrochloride Using Ferrous Sulphate in DMF Under Reflux Condition. *Asian Journal of Chemistry*, 24.3 (2012) 1401.
- [27] D.D. Patil, D.K. Mhaske, G.C. Wadhawa, Green Synthesis of 3,4-dihydropyrimidinone using ferrous sulphate as recyclable catalyst, *Journal of Pharmaceutical Research and Opinion*, 1(2011) 172-174.
- [28] S. S. Nayak, G. C. Wadhawa, K. B. Pathade, V. S. Shivankar, and N. A. Mirgane, Green synthesis of the plant assisted nanoparticles from Euphorbia neriifolia L. and its application in the degradation of dyes from industrial waste. *Plant Science Today*, 8.2 (2021) 380-385.
- [29] S.S. Nayak, G.C. Wadhawa, V.S. Shivankar, R. Inamadar, M.C. Sonawale, Phytochemical Analysis and Dpph Antioxidant Activity of Root And Bark of Syzygium Stocksii (Duthie) Plant, *Eur. J. Mol. Clin. Med.*, 7 (2021) 2655-2668
- [20] S.S. Nayak, G.C. Wadhawa, V.S. Shivankar, D.D. Patil, M.C. Sonawale, N.A. Mirgane, Tin oxide plant assisted nanoparticle catalyzed green synthesis of imidazole derivatives, *Materials Today: Proceedings*, 37 (2021) 2490-2494. <https://doi.org/10.1016/j.matpr.2020.08.301>.
- [31] G.C. Wadhawa, V.S. Shivankar, Y.A. Gaikwad, B.L. Ingale, B.R. Sharma, S. Hande, C.H. Gill, L.V. Gavali, One-pot synthesis of benzimidazoles the presence of lemon juice as a new and efficient catalyst. *European journal of pharmaceutical and medical research*, (ejpmr), 3(12) (2016) 556-558.
- [32] Wadhawa, G.C., Shivankar, V.S., Patil, S.S., Gaikwad, Y.A., Satere, A.V., Rode, B., Gill, C.H. and Gavali, L.V., Synthesis of 2-substituted benzimidazoles and 1, 5-disubstituted benzodiazepines using alum. *Rasayan J. Chem*, 10 (2017) 3-15.
- [33] G.C. Wadhawa, et al. Review on antibacterial activity of benzimidazole containing compounds. *World Journal of Pharmaceutical Research*, 7.7 (2018) 483-495.
- [34] D.D. Patil, G.C. Wadhawa. One pot synthesis of nitriles from aldehydes and hydroxylamine hydrochloride using sodium sulphate in DMF under reflux condition. *World Pharmacy and Pharmaceutical Sciences*, 3.1 (2013) 592-596.
- [35] D.D. Patil, D.K. Mhaske, G.C. Wadhawa. Antioxidant effect of the stem and leaves of Hibiscus esculentus Linn. *International Journal of Pharmaceutical Sciences and Research*, 2.6 (2011) 1464.

HOW TO CITE THIS ARTICLE

Pearl Rhea Fernandes, Ashwini Madnala, Nisha Donde, Amit Pawar, Maryappa sonawale, Green, Efficient

Synthesis of Baclofen, Ad. J. Chem. B, 4 (2022) 158-163.

DOI: 10.22034/ajcb.2022.342298.1120

URL: http://www.ajchem-b.com/article_152263.html